

The bipartite unconstrained 0-1 quadratic programming problem: polynomially solvable cases*

Abraham P. Punnen[†], Piyashat Sripratak[‡] and Daniel Karapetyan[§]

*Department of Mathematics, Simon Fraser University Surrey, Central City, 250-13450
102nd AV, Surrey, British Columbia, V3T 0A3, Canada*

Abstract

We consider the bipartite unconstrained 0-1 quadratic programming problem (BQP01) which is a generalization of the well studied unconstrained 0-1 quadratic programming problem (QP01). BQP01 has numerous applications and the problem is known to be MAX SNP hard. We show that if the rank of an associated $m \times n$ cost matrix $Q = (q_{ij})$ is fixed, then BQP01 can be solved in polynomial time. The complexity of this algorithm is quasi-polynomial if rank of Q is $O(\log^k n)$ for a fixed k . When Q is of rank one, we provide an $O(n \log n)$ algorithm and this complexity reduces to $O(n)$ with additional assumptions. Further, if $q_{ij} = a_i + b_j$ for some a_i and b_j , then BQP01 is shown to be solvable in $O(mn \log n)$ time. By restricting $m = O(\log n)$, we obtain yet another polynomially solvable case of BQP01 but the problem remains MAX SNP hard if $m = \sqrt[k]{n}$ for a fixed k . Finally, if the number of negative entries in Q is fixed, BQP01 is shown to be polynomially solvable whereas it is NP-hard if this number is $O(\sqrt[k]{n})$ for a fixed k . Many of the algorithms we obtained provide a unified framework for polynomially solvable cases of QP01 and BQP01.

Keywords: quadratic programming, 0-1 variables, polynomial algorithms, complexity, pseudo-Boolean programming.

1 Introduction

The unconstrained 0-1 quadratic programming problem (QP01) is to

$$\begin{aligned} &\text{Maximize } f(x) = x^T Q' x + c' x + c'_0 \\ &\text{Subject to } x \in \{0, 1\}^n, \end{aligned}$$

*This research work was supported by an NSERC Discovery grant awarded to Abraham P Punnen.

[†]apunnen@sfu.ca

[‡]psriprat@sfu.ca

[§]daniel.karapetyan@gmail.com

where Q' is an $n \times n$ matrix, c' is a row vector in R^n , and c'_0 is a constant. The problem QP01 is formulated and studied in various alternative formats. For example, without loss of generality, one can assume that c_0 is zero, c' is the zero vector and Q is symmetric. If c' is allowed to be arbitrary, then the diagonal elements of Q can be assumed to be zero, without loss of generality. In another variation, x is assumed to be in $\{-1, 1\}^n$ instead of $\{0, 1\}^n$. However, this variation can be reduced to the 0-1 version using a linear transformation. QP01 has been studied extensively in literature because of its various applications and rich theoretical structure. For further details and additional references on QP01 we refer to [5, 6, 9, 10, 14, 22]. The focus of this paper is on a problem closely related to QP01 which we call the *bipartite unconstrained 0-1 quadratic programming problem* (BQP01). The problem BQP01 can be defined as follows.

$$\begin{aligned} & \text{Maximize } f(x, y) = x^T Q y + c x + d y + c_0 \\ & \text{Subject to } x \in \{0, 1\}^m, y \in \{0, 1\}^n, \end{aligned}$$

where $Q = (q_{ij})$ is an $m \times n$ matrix, $c = (c_1, c_2, \dots, c_m)$ is a row vector in R^m , $d = (d_1, d_2, \dots, d_n)$ is a row vector in R^n and c_0 is a constant. Without loss of generality, we assume that $m \leq n$. Let

$$\bar{Q} = \left[\begin{array}{c|c} O_{m \times m} & Q \\ \hline O_{n \times m} & O_{n \times n} \end{array} \right], \bar{c} = [c \mid d] \text{ and } w^T = [x \mid y], \quad (1)$$

where $O_{n \times n}$, $O_{n \times m}$ and $O_{m \times m}$ are zero matrices. Then BQP01 can be formulated as the QP01

$$\begin{aligned} & \text{Maximize } f(w) = w^T \bar{Q} w + \bar{c} w + c_0 \\ & \text{Subject to } w \in \{0, 1\}^{m+n}. \end{aligned} \quad (2)$$

Thus, the exact and heuristic algorithms available to solve QP01 can be used directly to solve BQP01. However, such a transformation increases the problem size and could limit our ability in handling large scale problems. Further, BQP01 has additional structure that can be exploited to obtain efficient algorithms and derive interesting theoretical properties.

We can also show that QP01 is a special case of BQP01. Choose

$$Q = Q' + 2MI, \quad c = \frac{1}{2}c' - Me \text{ and } d = \frac{1}{2}c' - Me, \quad (4)$$

where I is an $n \times n$ identity matrix, $e \in R^n$ is an all one vector and M is a very large number. Then the resulting BQP01 solves QP01 since the penalty parameter M forces $x_i = y_i$ in an optimal solution of this modified BQP01. The transformation discussed in equation (4) is important since it provides additional flexibility in developing algorithms for QP01 through BQP01 formulations. This provides additional motivation to study BQP01.

BQP01 is also equivalent to the *unconstrained bilinear program*

$$\begin{aligned} \text{UBLP:} \quad & \text{Maximize } g(x, y) = x^T Q y + c x + d y + c_0 \\ & \text{Subject to } x \in [0, 1]^m, y \in [0, 1]^n, \end{aligned}$$

which is a special case of the *bilinear programming problem* (BLP) [3, 17, 32]. Note that the 0-1 variables in BQP01 are replaced by continuous variables in UBLP. It is well known that, for a BLP with variables $x^T = (x_1, x_2, \dots, x_m)$ and $y^T = (y_1, y_2, \dots, y_n)$, there exists an optimal solution (x^*, y^*) , where x^* is an extreme point of the constraint set defined by the x -variables and y^* is an extreme point of the constraint set defined by the y -variables [17]. In UBLP, these constraint sets are $\{0, 1\}$ -hypercubes and hence UBLP is equivalent to BQP01. Thus, BQP01 can also be solved using any algorithm for solving the BLP. However, by exploiting the special structure of UBLP, more efficient algorithms may be obtained. To the best of our knowledge, UBLP has not been studied in literature from the point of view of bilinear programs exploiting the inherent structure of the problem.

A graph theoretic interpretation of BQP01 can be given as follows. Let $I = \{1, 2, \dots, m\}$ and $J = \{1, 2, \dots, n\}$. Consider the bipartite graph $G = (I, J, E)$. For each node $i \in I$ and $j \in J$, respective costs c_i and d_j are prescribed. Further, for each $(i, j) \in E$, a cost q_{ij} is given. Then the *maximum weight induced subgraph problem* on G is to find $U_1 \subseteq I$, $U_2 \subseteq J$ such that $\sum_{i \in U_1} c_i + \sum_{j \in U_2} d_j + \sum_{(i,j) \in E^{1,2}} q_{ij}$ is maximized, where $E^{1,2}$ is the edge set of the subgraph of G induced $U_1 \cup U_2$. The maximum weight induced subgraph problem on G is precisely a BQP01, where $q_{ij} = 0$ if $(i, j) \notin E$.

Many well known combinatorial optimization problems can be modeled as BQP01. Consider the bipartite graph $G = (I, J, E)$ with $w_{ij} > 0$ being the weight of the edge $(i, j) \in E$. Then the *maximum weight biclique problem* (MWBP) [4, 30] is to find a biclique in G of maximum total edge-weight. Define

$$q_{ij} = \begin{cases} w_{ij} & \text{if } (i, j) \in E \\ -M & \text{otherwise,} \end{cases}$$

where M is a large positive number. Set c and d as zero vectors and also set $c_0 = 0$. Then the BQP01 with this choice of Q, c, d and c_0 solves the MWBP. This immediately shows that BQP01 is NP-hard and one can also establish some approximation hardness results with appropriate assumptions [4, 30]. MWBP has applications in data mining, clustering and bioinformatics [8, 31] which in turn become applications of BQP01.

Another application of BQP01 arises in approximating a matrix by a rank-one binary matrix [11, 20, 19, 23, 29]. For example, let $H = (h_{ij})$ be a given $m \times n$ matrix and we want to find an $m \times n$ matrix $A = (a_{ij})$, where $a_{ij} = u_i v_j$ and $u_i, v_j \in \{0, 1\}$, such that $\sum_{i=1}^m \sum_{j=1}^n (h_{ij} - u_i v_j)^2$ is minimized. The matrix A is called a rank one approximation of H and can be identified by solving the BQP01 with $q_{ij} = 1 - 2h_{ij}$, $c_i = 0$, $d_j = 0$ and $c_0 = 0$ for all $i \in I$ and $j \in J$. Binary matrix factorization is an important topic in mining discrete patterns in binary data [23, 29]. If u_i and v_j are required to be in $\{-1, 1\}$ then also the resulting approximation problem can be formulated as a BQP01. The maximum

cut problem on a bipartite graph can be formulated as a BQP01 and this gives yet another application of the model. BQP01 can also be used to find approximations to the cut-norm of a matrix [2]. When the matrix Q' is positive semidefinite, QP01 can be solved directly as a BQP01. This follows from a corresponding result from BLP [17] and the equivalence between UBLP and BQP01. The applications and potential theoretical developments make it interesting to investigate BQP01 further.

To the best of our knowledge, BQP01 has not been thoroughly investigated in literature, especially from the point of view of polynomially solvable special cases. The primary focus of this work is to identify such special cases and provide a concise review of the available results from the literature on BQP01.

We show that BQP01 can be solved in polynomial time if the rank of Q is fixed. It may be noted that the corresponding version of QP01 is NP-hard even if Q is of rank one [13]. With the additional restrictions that Q' is positive semidefinite and $c' = 0$, QP01 can be solved in polynomial time [1, 10, 16, 24]. Our algorithm can also be used to solve this special case of QP01 in polynomial time with the additional flexibility that c can be an arbitrary vector. When Q is of rank one, we show that BQP01 can be solved in $O(n \log n)$ time. In addition, we obtain an $O(n)$ algorithm for a special case of this rank-one problem. When $q_{ij} = a_i + b_j$, we present an $O(mn \log n)$ algorithm. Further, we observe that if $m = O(\log n)$ then BQP01 can be solved in polynomial time but if $m = \sqrt[k]{n}$ for a fixed k , the problem remains MAX SNP hard. If $m = O(\log^k n)$ then BQP01 can be solved in quasi-polynomial time. Similarly, if the number of negative entries in Q are $O(\log n)$ then BQP01 is solvable in polynomial time but if the number of negative entries are $O(\sqrt[k]{n})$ for a fixed k , the problem becomes NP-hard.

The paper is organized as follows. In Section 2, we present various equivalent formulations of BQP01. Section 3 deals with polynomially solvable special cases. Concluding remarks are given Section 4.

2 Equivalent formulations

QP01 is often presented in literature in various equivalent forms. Similar equivalent formulations can be obtained for BQP01 as well which are summarized in this section. Although many of the transformations discussed here follows almost directly from the corresponding transformation for QP01, we present them here for completeness, comparison, and for notational convenience. Further, these transformations, albeit simple, preserve the bipartite structure of the problem which allows us establishing interesting complexity results.

When the linear and constant terms from BQP01 are dropped, we get the homogeneous version of the problem defined as,

$$\begin{aligned} \text{BQP01(H):} \quad & \text{Maximize } f(x, y) = x^T Q y \\ & \text{Subject to } x \in \{0, 1\}^m, y \in \{0, 1\}^n. \end{aligned}$$

In fact, the general BQP01 can be represented in the homogeneous form by introducing two

additional variables. Let

$$\bar{Q} = \left[\begin{array}{c|c} Q & c^T \\ \hline d & c_0 \end{array} \right]. \quad (5)$$

Then BQP01 is equivalent to

$$\begin{aligned} \text{BQP01}(\bar{H}): \quad & \text{Maximize } f(\bar{x}, \bar{y}) = \bar{x}^T \bar{Q} \bar{y} \\ & \text{Subject to } \bar{x} \in \{0, 1\}^{m+1}, \bar{y} \in \{0, 1\}^{n+1}, \bar{x}_{m+1} = 1, \bar{y}_{n+1} = 1, \end{aligned}$$

where $\bar{x}_{m+1}$ and $\bar{y}_{n+1}$ are respectively the $(m+1)$ th and $(n+1)$ th components of $\bar{x}$ and $\bar{y}$. The restrictions $\bar{x}_{m+1} = 1, \bar{y}_{n+1} = 1$ in BQP01($\bar{H}$) can be dropped by replacing c_0 with a large number M in $\bar{Q}$. Thus, we have

Remark 1. *A BQP01 can always be represented in an equivalent homogeneous form in the sense that the two problems have identical optimal solutions set.*

Another important variation of BQP01 is obtained by restricting the variables to 1 or -1 instead of 0 or 1. We call this the *cut version* of BQP01 and denote it by BQP-11. The problem BQP-11 can be stated as

$$\begin{aligned} \text{BQP-11}: \quad & \text{Maximize } \phi(x, y) = x^T Q y + cx + dy + c_0 \\ & \text{Subject to } x \in \{-1, 1\}^m, y \in \{-1, 1\}^n. \end{aligned}$$

By dropping the linear and constant terms from BQP-11, we get the corresponding homogeneous version

$$\begin{aligned} \text{BQP-11(H)}: \quad & \text{Maximize } \phi(x, y) = x^T Q y \\ & \text{Subject to } x \in \{-1, 1\}^m, y \in \{-1, 1\}^n. \end{aligned}$$

As in the case of BQP01, its cut version BQP-11 can also be represented in the corresponding homogeneous form by introducing two additional variables. Let $\bar{Q}$ be defined as in equation (5) and $\bar{x} \in \{-1, 1\}^{m+1}, \bar{y} \in \{-1, 1\}^{n+1}$. Then BQP-11 is equivalent to

$$\begin{aligned} \text{BQP-11}(\bar{H}): \quad & \text{Maximize } \phi(\bar{x}, \bar{y}) = \bar{x}^T \bar{Q} \bar{y} \\ & \text{Subject to } \bar{x} \in \{-1, 1\}^{m+1}, \bar{y} \in \{-1, 1\}^{n+1}, \bar{x}_{m+1} = 1, \bar{y}_{n+1} = 1. \end{aligned}$$

Note that the restriction $\bar{x}_{m+1} = 1, \bar{y}_{n+1} = 1$ in BQP-11($\bar{H}$) can be replaced by $\bar{x}_{m+1} = \bar{y}_{n+1}$ without affecting the optimal solution of BQP-11($\bar{H}$). This is possible because, for any solution $\bar{x}, \bar{y}$ with $\bar{x}_{m+1} = -1, \bar{y}_{n+1} = -1$, the solution $-\bar{x}, -\bar{y}$ satisfies $-\bar{x}_{m+1} = 1, -\bar{y}_{n+1} = 1$ and $\phi(\bar{x}, \bar{y}) = \phi(-\bar{x}, -\bar{y})$. Further, the condition $\bar{x}_{m+1} = \bar{y}_{n+1}$ can be dropped by replacing c_0 with a large number M in $\bar{Q}$ resulting in a homogeneous version of BQP-11. Thus, we have

Remark 2. *A BQP-11 can always be represented in an equivalent homogeneous version in the sense that the two problems have identical optimal solutions set.*

Consider the linear transformation

$$x = 2w - e_m \text{ and } y = 2z - e_n, \quad (6)$$

where e_m and e_n are all-one vectors in R^m and R^n , respectively. Using (6), BQP-11 can be reduced to the BQP01

$$\begin{aligned} & \text{Maximize } f(w, z) = w^T \tilde{Q}z + \tilde{c}w + \tilde{d}z + \tilde{c}_0 \\ & \text{Subject to } w \in \{0, 1\}^m, z \in \{0, 1\}^n, \end{aligned}$$

where $\tilde{Q} = 4Q$, $\tilde{c} = 2(c - (Qe_n)^T)$, $\tilde{d} = 2(d - e_m^T Q)$ and $\tilde{c}_0 = e_m^T Qe_n - ce_m - de_n + c_0$.

Similarly, using the linear transformation

$$x = \frac{1}{2}(w + e_m) \text{ and } y = \frac{1}{2}(z + e_n) \quad (7)$$

BQP01 can be reduced to the BQP-11

$$\begin{aligned} & \text{Maximize } \phi(w, z) = w^T \hat{Q}z + \hat{c}w + \hat{d}z + \hat{c}_0 \\ & \text{Subject to } w \in \{-1, 1\}^m, z \in \{-1, 1\}^n \end{aligned}$$

where $\hat{Q} = \frac{1}{4}Q$, $\hat{c} = \frac{1}{4}(Qe_n)^T + \frac{1}{2}c$, $\hat{d} = \frac{1}{4}e_m^T Q + \frac{1}{2}d$ and $\hat{c}_0 = \frac{1}{4}e_m^T Qe_n + \frac{1}{2}ce_m + \frac{1}{2}de_n + c_0$.

It is well known that the maximum weight cut problem (MaxCut) on a general graph G is equivalent to QP01 [14]. In the maximum weight cut problem, if we restrict the graph G to be bipartite, we get an instance of the *bipartite maximum weight cut problem* which is denoted by B-MaxCut. Indeed, viewing B-MaxCut as a general MaxCut problem yields an equivalent QP01. Many approximation algorithms for MaxCut assume that the associated edge weights are nonnegative [12]. However, for nonnegative edge weights, B-MaxCut is a trivial problem since the generic bipartition of the underlying graph gives the optimal cut. Thus, the topic that is more interesting is when the edge weights take positive as well as negative values. Developing approximation algorithms for BQP01 or for B-MaxCut is not the focus of this paper. However, we do want to indicate the equivalence between B-MaxCut and BQP01 to further understand the inherent complexity of the problems.

Let $G = (I, J, E)$ be a bipartite graph. Two vectors $x \in \{-1, 1\}^m$ and $y \in \{-1, 1\}^n$ determine a cut $(U_1 \cup U_2, H_1 \cup H_2)$ in G if and only if $U_1 = \{i \in I : x_i = 1\}$, $H_1 = \{i \in I : x_i = -1\}$, $U_2 = \{j \in J : y_j = 1\}$, and $H_2 = \{j \in J : y_j = -1\}$. We call (x, y) the *incidence vector* of the cut $(U_1 \cup U_2, H_1 \cup H_2)$. Let q_{ij} be the weight of the edge (i, j) in G . Then the value of the cut $(U_1 \cup U_2, H_1 \cup H_2)$ is given by

$$\sum_{i \in U_1, j \in H_2} q_{ij} + \sum_{i \in H_1, j \in U_2} q_{ij} = \sum_{x_i = -y_j} q_{ij} \quad (8)$$

Theorem 3. *The problems BQP01 and B-MaxCut are equivalent in the sense that:*

1. For any instance of BQP01, it is possible to construct a complete bipartite graph G such that an optimal solution to the B-MaxCut problem on G gives an optimal solution to BQP01.
2. For any instance of B-MaxCut on a bipartite graph $G = (I, J, E)$, it is possible to construct an instance of BQP01 with an $m \times n$ cost matrix Q such that an optimal solution to the BQP01 gives an optimal solution to the B-MaxCut problem on G .

Proof. Since BQP01 is equivalent to a BQP-11(H) (see remarks 1 and 2), without loss of generality we assume that BQP01 is given in the equivalent homogeneous cut form BQP-11(H). Now,

$$\phi(x, y) = \sum_{ij} q_{ij} x_i y_j = \sum_{x_i=y_j} q_{ij} - \sum_{x_i=-y_j} q_{ij} = \sum_{ij} q_{ij} - 2 \sum_{x_i=-y_j} q_{ij}$$

Since $\sum_{ij} q_{ij}$ is a constant, maximizing $\phi(x, y)$ is equivalent to maximizing $-2 \sum_{\bar{x}_i=-\bar{y}_j} q_{ij}$. Thus, by solving the B-MaxCut problem on a complete bipartite graph $K_{m,n}$ with $-2q_{ij}$ chosen as the weight of the edge (i, j) , we can solve BQP-11(H).

To establish the second part of the theorem, we show that the B-MaxCut problem on G with edge weights c_{ij} for $(i, j) \in E$ can be solved as a BQP-11(H). Let $(U_1 \cup U_2, H_1 \cup H_2)$ be a cut in G and (x, y) be the corresponding incidence vector. Let $g(U, H) = \sum_{x_i=-y_j} c_{ij}$ be the value of the cut $(U_1 \cup U_2, H_1 \cup H_2)$. Then it can be verified that $g(U, H) = \frac{1}{2} \sum_{ij} c_{ij} - \frac{1}{2} \sum_{ij} c_{ij} x_i y_j$ and, hence, maximizing $g(U, H)$ is equivalent to solving a BQP-11(H) with cost matrix $q_{ij} = -\frac{1}{2} c_{ij}$. $\square$

3 Polynomially solvable cases

Alon and Naor [2] showed that BQP-11(H) is MAX SNP hard. Using the linear transformation (6), BQP-11(H) can be reduced to a BQP01 and the reduction preserves the objective function values of the corresponding solutions. Thus, BQP01 is also MAX SNP hard. Now let us focus on polynomially solvable special cases of BQP01.

Polynomially solvable special cases of QP01 have been investigated by many authors [1, 7, 10, 16, 21, 24, 27]. Since BQP01 can be formulated as a QP01 using the transformation (3), whenever the transformed data satisfies the known conditions for polynomial solvability of QP01, we can solve the corresponding BQP01 also in polynomial time. So let us focus on new polynomially solvable cases that exploit the special structure of BQP01.

Theorem 4. *BQP01 can be solved in polynomial time if $m = O(\log n)$ but remains MAX SNP hard if $m = O(\sqrt[k]{n})$ for any constant k . Further, if m is fixed, BQP01 can be solved in $O(n)$ time.*

Proof. Consider a 0-1 m -vector x^0 set $x = x^0$. Construct a vector y^0 as follows:

$$y_j^0 = \begin{cases} 1 & \text{if } \sum_{i=1}^m q_{ij} x_i^0 + d_j > 0 \\ 0 & \text{otherwise} \end{cases} \quad (9)$$

When x is restricted to x^0 , the choice of y^0 given by equation (9) yields the maximum objective function value. Thus, y^0 and the objective function value of the solution (x^0, y^0) can be identified in $O(mn)$ time. Repeat this for all possible choices of the vector x^0 , compute the corresponding y^0 , and choose the overall best solution obtained in this process. This will be an optimal solution to BQP01. Since the number of different choices for x^0 is 2^m and by our assumption $m = k \log n$ for some constant k , BQP01 can be solved in $O(mn2^m) = O(n^{k+1} \log n)$ time. If m is fixed, the complexity reduces to $O(n)$.

To establish that BQP01 is MAX SNP hard when $m = O(\sqrt[k]{n})$, consider an instance of BQP01 with data Q, c, d and c_0 . Let

$$\bar{Q} = \left[\begin{array}{c|c} Q & O_{m \times n^k} \\ \hline O_{(n-m) \times (n-m)} & O_{(n-m) \times n^k} \end{array} \right], \bar{c} = [c \mid \tilde{c}], \text{ and } \bar{d} = [d \mid \tilde{d}],$$

where $O_{m \times n^k}$, $O_{(n-m) \times (n-m)}$ and $O_{(n-m) \times n^k}$ are zero matrices, $\tilde{c}$ is a zero vector of dimension $n - m$ and $\tilde{d}$ is a zero vector of dimension n^k . Now consider the instance of BQP01 with data $\bar{Q}, \bar{c}, \bar{d}$ and c_0 . It can be verified that this new instance of BQP01 satisfies the condition of the theorem. Further, for every solution to the original BQP01, we can find a solution to the constructed instance of BQP01, where the corresponding objective function values are identical and viceversa. Since BQP01 is MAX SNP hard, the result follows. $\square$

If $m = O(\log^k n)$ for fixed k , the procedure discussed in the proof of Theorem 4 becomes quasi-polynomial. Thus, BQP01 can be solved in quasi-polynomial time if $m = O(\log^k n)$ for fixed k . Let us now consider a generalization of Theorem 4 where rank of Q is fixed, instead of the number of rows of Q .

QP01 is polynomially solvable if Q' is a symmetric positive semidefinite matrix [1, 10, 16, 24] with fixed rank and $c' = 0$. If c' is allowed to have arbitrary elements, the problem is NP-hard even if the rank of Q' is one [13]. We now show that BQP01 is solvable in polynomial time if rank of Q is fixed. No assumption is made on any other property of Q such as positive semidefiniteness and no restrictions on c and d are imposed. As a byproduct of this result, we have an alternative polynomial time algorithm to solve QP01 when rank of Q' is fixed and Q' is positive semidefinite with the additional flexibility that the vector c' could be arbitrary. First, let us prove some preliminary results.

Consider the multiparametric linear programming problem (MLP)

$$\begin{aligned} f(\lambda) &= \max cx \\ \text{Subject to: } Ax &= \lambda \\ x &\in [0, 1]^m, \end{aligned}$$

where A is a $p \times m$ matrix of full row rank and $\lambda^T = (\lambda_1, \lambda_2, \dots, \lambda_p) \in R^p$. The j th column of the matrix A is denoted by A_j . A partition $(\mathcal{B}, \mathcal{L}, \mathcal{U})$ of $I = \{1, 2, \dots, m\}$ with $|\mathcal{B}| = p$ is referred to as a *basis structure* for MLP. For each basic feasible solution of MLP, a basis structure $(\mathcal{B}, \mathcal{L}, \mathcal{U})$ is associated, where $\mathcal{L}$ is the index set of nonbasic variables at the lower bound 0, $\mathcal{U}$ is the index set of non-basic variables at the upper bound 1 and

$\mathcal{B} = \{B_1, B_2, \dots, B_p\}$ is the index set of basic variables. Let B be the $p \times p$ matrix with its i th column A_{B_i} . Then the set $\mathcal{B}$ is called a basis set or simply a *basis* and B is the associated *basis matrix*. A basis provides an implicit ordering of its elements. Thus, x_{B_i} is called the i th basic variable with respect to $\mathcal{B}$. Let $C_B = (c_{B_1}, c_{B_2}, \dots, c_{B_p})$. Then the basis structure $(\mathcal{B}, \mathcal{L}, \mathcal{U})$ is dual feasible if and only if [26]

$$C_B B^{-1} A_j - c_j \geq 0 \text{ for } j \in \mathcal{L} \quad (10)$$

$$C_B B^{-1} A_j - c_j \leq 0 \text{ for } j \in \mathcal{U}. \quad (11)$$

Conditions (10) and (11) are also known as *reduced cost optimality conditions* [26]. Let $(\mathcal{B}, \mathcal{L}, \mathcal{U})$ be a dual feasible basis structure for MLP. Then the basic solution of MLP corresponding to this basis structure is optimal for all $\lambda \in R^p$ satisfying

$$\mathbf{0} \leq B^{-1} \lambda - B^{-1} A^u e \leq \mathbf{1}, \quad (12)$$

where A^u is the submatrix formed by columns of A corresponding to the indices of $\mathcal{U}$, e is vector of size $|\mathcal{U}|$ with all entry equal to 1, $\mathbf{0}$ is the zero vector in R^p and $\mathbf{1}$ is the vector in R^p with all entries equal to 1. The polyhedral set represented by (12) is called the *characteristic region* of the basis structure $(\mathcal{B}, \mathcal{L}, \mathcal{U})$. A dual feasible basis structure $(\mathcal{B}, \mathcal{L}, \mathcal{U})$ is *dual non-degenerate* if (10) and (11) are satisfied as strict inequalities. If any of these inequalities is satisfied as an equality, then $(\mathcal{B}, \mathcal{L}, \mathcal{U})$ is a *dual degenerate* basis structure.

Let S be the collection of all the extreme points of the characteristic regions associated with all dual feasible basis structures of MLP.

Lemma 5. $|S| \leq {}^m\mathcal{C}_p 2^p$.

Proof. For simplicity, we assume that all dual feasible basis structures of MLP are dual non-degenerate. This is not a restriction since we can achieve this by an ϵ -perturbation of the cost vector c in MLP by appropriately small ϵ and this change will not underestimate $|S|$. Thus, the inequalities (10) and (11) are satisfied as strict inequalities. Consequently, given $\mathcal{B}$, the choice of $\mathcal{L}$ and $\mathcal{U}$ is unique for any dual feasible basis structure, i.e., if $(\mathcal{B}, \mathcal{L}_1, \mathcal{U}_1)$ and $(\mathcal{B}, \mathcal{L}_2, \mathcal{U}_2)$ are two dual non-degenerate basis structures for MLP then at most one of them can be dual feasible. Thus, there exist at most ${}^m\mathcal{C}_p$ dual feasible and dual non-degenerate basis structures for MLP. The characteristic region of such a basis structure $(\mathcal{B}, \mathcal{L}, \mathcal{U})$ is defined by the inequalities (12). Each extreme point of this polyhedron is determined by the unique solution of p tight inequalities from (12). Since there are at most 2^p choices for these tight inequalities, the result follows. $\square$

Since $f(\lambda)$ is a piecewise linear concave function [15], it also defines a polyhedral set and the extreme points of this set are precisely elements of S . These extreme points are sometimes called *breakpoints* of $f(\lambda)$. Thus, $f(\lambda)$ will have at most ${}^m\mathcal{C}_p 2^p$ breakpoints.

Theorem 6. *If $x(\lambda^0)$ is an optimal basic feasible solution for MLP at $\lambda = \lambda^0 \in S$ then $x(\lambda^0) \in \{0, 1\}^m$.*

Proof. Let $(\mathcal{B}, \mathcal{L}, \mathcal{U})$ be a basis structure that generated the extreme point λ^0 and B be the associated basis matrix. Then the vector of basic variables x_B associated with $(\mathcal{B}, \mathcal{L}, \mathcal{U})$ is given by

$$x_B = B^{-1}\lambda^0 - B^{-1}A^ue. \quad (13)$$

But λ^0 is an extreme point of $f(\lambda)$. Thus from equation (12), λ^0 is the solution of the system $B^{-1}\lambda^0 - B^{-1}A^ue = \tau$ for some 0-1 vector τ . Thus from (13), x_B is a 0-1 vector. The non-basic variables of $x(\lambda)$ by definition take 0-1 values and the result follows. $\square$

Let us now consider BQP01 where rank of Q is p , a fixed number. Since BQP01 is equivalent to UBLP, this problem can be stated as

$$\begin{aligned} \text{UBLP}(p): \quad & \text{Maximize } \sum_{k=1}^p a^k x b^k y + cx + dy \\ & \text{Subject to: } x \in [0, 1]^m, y \in [0, 1]^n, \end{aligned}$$

where $a^k = (a_1^k, a_2^k, \dots, a_m^k)$ and $b^k = (b_1^k, b_2^k, \dots, b_n^k)$ for $k = 1, 2, \dots, p$. Consider the multi-parametric linear program (MLP1) [15]

$$\begin{aligned} \text{MLP1:} \quad & h_1(\lambda) = \max cx \\ & \text{Subject to: } a^k x = \lambda_k \text{ for } k = 1, 2, \dots, p \\ & x \in [0, 1]^m, \end{aligned}$$

where $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_p)$. Then $h_1(\lambda)$ is a piecewise linear concave function [15]. Let S_1 be the set of breakpoints of $h_1(\lambda)$ and $x(\lambda)$ be an optimal basic feasible solution of MLP1 at $\lambda \in S_1$. By Theorem 6, $x(\lambda) \in \{0, 1\}^m$. Let $y(\lambda) = (y_1(\lambda), y_2(\lambda), \dots, y_n(\lambda))$ be an optimal solution to UBLP(p) when x is restricted to $x(\lambda)$. Then it can be verified that $y(\lambda)$ satisfies

$$y_j(\lambda) = \begin{cases} 1 & \text{if } d_j + \sum_{k=1}^p b_j^k \sum_{i \in I} a_i^k x_i(\lambda) > 0 \\ 0 & \text{otherwise} . \end{cases} \quad (14)$$

Theorem 7. *There exists an optimal solution to UBLP(p) amongst the solutions $\{x(\lambda), y(\lambda) : \lambda \in S_1\}$.*

Proof. UBLP(p) is equivalent to the bilinear program

$$\begin{aligned} \text{BLP1:} \quad & \text{Maximize } \sum_{k=1}^p \lambda_k b^k y + cx + dy \\ & \text{Subject to: } a^k x = \lambda_k \text{ for } k = 1, 2, \dots, p \\ & x \in [0, 1]^m, y \in [0, 1]^n. \end{aligned}$$

Let $h(\lambda)$ be the optimal objective function value of BLP1 where λ is fixed. Now, $h(\lambda)$ can be decomposed into $h_1(\lambda) + h_2(\lambda)$, where

$$h_2(\lambda) = \max \sum_{k=1}^p \lambda_k b^k y + dy$$

Subject to: $y \in [0, 1]^n$,

and $h_1(\lambda)$ is as defined in MLP1. Then the optimal objective function value of UBLP(p) can be identified by finding the global maximum of $h(\lambda)$ over all $\lambda \in R^p$ for which BLP1 is feasible. As a function of λ , $h_2(\lambda)$ is piecewise linear convex [15] and $h_1(\lambda)$ is piecewise linear concave [15]. Thus, $h(\lambda)$ is piecewise linear but need not be convex or concave. But $h_1(\lambda)$ is linear when λ is restricted to its characteristic region. Thus, $h(\lambda)$ is convex when λ is restricted to any characteristic region associated with $h_1(\lambda)$. Hence, the maximum of $h(\lambda)$ is attained at an extreme point of $h_1(\lambda)$. Let $\lambda = \lambda^0$ be such an extreme point and $x(\lambda^0)$ be an optimal basic feasible solution of MLP1 at $\lambda = \lambda^0$. By Theorem 6, $x(\lambda^0)$ is a 0-1 solution and it yields the optimal value $y(\lambda)$ as given by (14) for the y variables for UBLP(p) subject to the condition that $x = x^0(\lambda)$. Since one of the $x(\lambda)$ for $\lambda \in S$ gives an optimal x -value, the result follows. $\square$

Based on Theorem 7, UBLP(p) and hence BQP01 can be solved by generating the set S_1 , computing the set $S_2 = \{x(\lambda), y(\lambda) : \text{for all } \lambda \in S_1\}$ and choosing the best solution in S_2 . Since a dense matrix inversion of size p can be carried out in $O(p^3)$ time, it can be verified that the overall complexity of this process is $O(m\mathcal{C}_p 2^p (p^3 + mp + n)) = O\left(\left(\frac{me}{p}\right)^p 2^p (p^3 + mp + n)\right)$. Since UBQP(p) and BQP01 are equivalent when Q is of rank p we have,

Theorem 8. *BQP01 can be solved in $O\left(\left(\frac{m}{p}\right)^p e^p (p^3 + mp + n)\right)$ time when rank of $Q = p$.*

Note that for a fixed p the above bound is polynomial and for $p = O(\log n)$ it is quasipolynomial.

For specific choices of p , the complexity of this procedure may be improved. We illustrate this for the case when $p = 1$. Theorem 8 guarantees that if rank of Q is 1, BQP01 can be solved in $O(mn)$ time. We now show that the problem can be solved in $O(n \log n)$ time by careful organization of our computations. As in the general case, let us consider the bilinear equivalent version:

$$\begin{aligned} \text{UBLP}(1): \quad & \text{Maximize } axby + cx + dy \\ & \text{Subject to: } x \in [0, 1]^m, y \in [0, 1]^n, \end{aligned}$$

where $a = (a_1, a_2, \dots, a_m)$, $c = (c_1, c_2, \dots, c_m) \in R^m$ and $b = (b_1, b_2, \dots, b_n)$, $d = (d_1, d_2, \dots, d_n) \in R^n$. Let $A^- = \{i : a_i < 0\}$, $A^0 = \{i : a_i = 0\}$ and $A^+ = \{i : a_i > 0\}$. Define $\underline{\lambda} = \sum_{i \in A^-} a_i$ and $\bar{\lambda} = \sum_{i \in A^+} a_i$, where summation over the empty set is taken as zero. Note that $\underline{\lambda}$ and

$\bar{\lambda}$ are respectively the smallest and the largest values of ax when $x \in [0, 1]^m$. Consider the *parametric continuous knapsack problem* (PKP(λ)) given below.

$$\begin{aligned} \text{PKP}(\lambda): \quad & \text{Maximize } cx \\ & \text{Subject to: } ax = \lambda \\ & x \in [0, 1]^m, \text{ and } \underline{\lambda} \leq \lambda \leq \bar{\lambda}. \end{aligned}$$

This is a special case of MLP1 for $p = 1$. Let $h_1(\lambda)$ be the optimal objective function value of PKP(λ) for a given λ . Then for $\underline{\lambda} \leq \lambda \leq \bar{\lambda}$, $h_1(\lambda)$ is a piecewise linear concave function [26]. Let $\underline{\lambda} = \lambda_1 < \lambda_2 < \dots < \lambda_r = \bar{\lambda}$ be the breakpoints of $h_1(\lambda)$, x^k be an optimal basic feasible solution of PKP(λ) for $\lambda \in [\lambda_k, \lambda_{k+1}]$, $1 \leq k \leq r-1$, and x^r be an optimal basic feasible solution to PKP(λ^r). Let y^k be an optimal solution to UBLP(1) when x is restricted to x^k , the vector y^k can be identified by appropriate modification of the equation (14). By Theorem 7, there exists an optimal solution to UBLP(1) amongst the solutions $x^k, y^k : k = 1, 2, \dots, r$.

From Lemma 5, the number of breakpoints of $h_1(\lambda)$ is at most $2m$. We now observe that the number of breakpoints of $h_1(\lambda)$ cannot be more than $m+1$ and obtain closed form values of these breakpoints. Define

$$\alpha_i = \begin{cases} c_i & \text{if } a_i \geq 0 \\ -c_i & \text{if } a_i < 0. \end{cases}$$

Let $T = \left\{ \frac{\alpha_i}{|a_i|} : i = 1, 2, \dots, m, a_i \neq 0 \right\}$ and consider a descending arrangement

$$\frac{\alpha_{\pi_1}}{|a_{\pi_1}|} > \frac{\alpha_{\pi_2}}{|a_{\pi_2}|} > \dots > \frac{\alpha_{\pi_r}}{|a_{\pi_r}|} \quad (15)$$

of all distinct elements of T . Let $T(k) = \left\{ i : \frac{\alpha_{\pi_k}}{|a_{\pi_k}|} = \frac{\alpha_i}{|a_i|} \right\}$. Then the breakpoints of $h_1(\lambda)$ are given by

$$\lambda_1 = \underline{\lambda} \text{ and } \lambda_k = \lambda_{k-1} + \sum_{i \in T(k)} |a_i| \text{ for } k = 2, 3, \dots, r.$$

An optimal solution to PKP(λ) at $\lambda = \lambda_k$ for $k = 1, 2, \dots, r$ can be identified recursively as

$$x_i^1 = \begin{cases} 1 & \text{if } a_i = 0 \text{ and } c_i > 0 \text{ or } a_i < 0 \\ 0 & \text{otherwise} \end{cases}$$

$$\text{and } x_i^k = \begin{cases} x_i^{k-1} & \text{if } i \notin T(k) \\ 1 & \text{if } i \in T(k) \text{ and } a_i > 0 \\ 0 & \text{otherwise,} \end{cases}$$

with the corresponding objective function values

$$h_1(\lambda_1) = \sum_{i=1}^m c_i x_i^1 \text{ and } h_1(\lambda_k) = h_1(\lambda_{k-1}) + \sum_{i \in T(k)} \alpha_i.$$

Given $h(\lambda_k)$ and x^k , $h(\lambda_{k+1})$ and x^{k+1} can be identified in $O(|T(k+1)|)$ time. The complexity for generating these solutions and breakpoints are dominated by that of constructing the descending arrangement (15) which is $O(m \log m)$. Note that $h_1(\lambda)$ has at most $m+1$ breakpoints and given x^k a corresponding solution y^k can be computed in $O(n)$ time. This leads to a complexity of $O(mn)$. The bottleneck operation here is the computation of y^k for $k = 1, 2, \dots, r$. We now show that these points can be identified in $O(n \log n)$ time.

Consider the *parametric unconstrained linear optimization problem*

$$\begin{aligned} \text{ULP}(\mu): \quad & \text{Maximize } dy + \mu by \\ & \text{Subject to: } y \in [0, 1]^m \text{ and } \underline{\lambda} \leq \mu \leq \bar{\lambda}. \end{aligned}$$

Let $h_2(\mu)$ be the optimal objective function value of $\text{ULP}(\mu)$. Then $h_2(\mu)$ is a piecewise linear convex function.

Let $S^+ = \{j : d_j + \underline{\lambda} b_j \geq 0\}$, $S^- = \{j : d_j + \underline{\lambda} b_j < 0\}$, $B^+ = \{j : b_j > 0\}$, and $B^- = \{j : b_j < 0\}$. Also, let $T_2 = \{-\frac{d_j}{b_j} : j \in (S^+ \cap B^-) \cup (S^- \cap B^+)\}$ and consider an ascending arrangement

$$\underline{\lambda} < -\frac{d_{\tau_1}}{b_{\tau_1}} < -\frac{d_{\tau_2}}{b_{\tau_2}} \dots < -\frac{d_{\tau_t}}{b_{\tau_t}} \quad (16)$$

of all distinct elements of T_2 greater than $\underline{\lambda}$. Let $\mu_1, \mu_2, \dots, \mu_t$ be the breakpoints of $h_2(\mu)$. Then

$$\mu_\ell = -\frac{d_{\tau_\ell}}{b_{\tau_\ell}} \text{ for } \ell = 1, 2, \dots, t.$$

Let $\Delta_i = \{j : j \in (S^+ \cap B^-) \cup (S^- \cap B^+) \text{ and } \frac{d_j}{b_j} = \frac{d_{\tau_i}}{b_{\tau_i}}\}$. Then the optimal solution y_ℓ for $\text{ULP}(\mu)$ corresponding to the breakpoint μ_ℓ for $\ell = 1, 2, \dots, t$ is given recursively by

$$y_j^\ell = \begin{cases} y_j^{\ell-1} & \text{if } j \notin \Delta_\ell \\ 1 & \text{if } j \in \Delta_\ell \text{ and } y_j^{\ell-1} = 0 \\ 0 & \text{if } j \in \Delta_\ell \text{ and } y_j^{\ell-1} = 1. \end{cases}$$




Figure 1: An example of $h_1(\lambda)$ and $h_2(\mu)$ when $a = [2, 2, -3, 4, -2]$, $c = [4, 5, 6, 10, 5]$, $b = [1, 1, -4, 0, -1, -2, 1]$, $d = [5, -2, 3, 3, 4, 0, 2]$.

where

$$y_j^0 = \begin{cases} 1 & \text{if } j \in S^+ \\ 0 & \text{otherwise} \end{cases}.$$

Define

$$\begin{aligned} D^0 &= \sum_{j \in S^+} d_j, \quad B^0 = \sum_{j \in S^+} b_j, \\ D^\ell &= D^{\ell-1} - \sum_{j \in \Delta_\ell, y_j^{\ell-1}=1} d_j + \sum_{j \in \Delta_\ell, y_j^{\ell-1}=0} d_j \text{ and} \\ B^\ell &= B^{\ell-1} - \sum_{j \in \Delta_\ell, y_j^{\ell-1}=1} b_j + \sum_{j \in \Delta_\ell, y_j^{\ell-1}=0} b_j. \end{aligned}$$

Then the optimal objective function value at μ_ℓ is given by $h_2(\mu_\ell) = D^\ell + \mu_\ell B^\ell$.

Given $y^{\ell-1}$, $D^{\ell-1}$ and $B^{\ell-1}$, we can compute y^ℓ , D^ℓ , and B^ℓ in $O(|\Delta_\ell|)$ time and, hence, $h_2(\mu_\ell)$ and y^ℓ can be identified in $O(|\Delta_\ell|)$ time. Since $\Delta_\ell \cap \Delta_k = \emptyset$ for $\ell \neq k$, y^ℓ and $h_2(\mu^\ell)$ for $\ell = 1, 2, \dots, t$ can be identified in $O(n \log n)$ time.

Now the algorithm for solving UBLP(1) can be described as follows. First, compute $x^1, y^1, h_1(\underline{\lambda})$ and $h_2(\underline{\lambda})$. Set $f(x^1, y^1) = h_1(\underline{\lambda}) + h_2(\underline{\lambda})$. Sort all breakpoints of $h_1(\lambda)$ and

$h_2(\mu)$ for $\underline{\lambda} \leq \lambda, \mu \leq \bar{\lambda}$ (see Fig: 1) and scan these breakpoints starting from $\underline{\lambda}$ in the increasing order. As we pass breakpoints of $h_2(\mu)$ keep updating the solution y of $\text{ULP}(\mu)$ corresponding to this breakpoint and the objective function value of this solution until we hit a breakpoint λ_k of $h_1(\lambda)$. At this point compute the solution x^k and $h_1(\lambda_k)$. The most recent solution y identified is selected as y^k and compute $h_2(\lambda_k)$. Note that $h_2(\lambda_k)$ can be obtained in $O(1)$ time using slope of $h_2(\mu)$ for the interval containing λ^k . Update $f(x^k, y^k)$ and the process is continued until all breakpoints of $h_1(\lambda)$ including $\bar{\lambda}$ are examined and the overall best solution is selected. It is not difficult to verify that the complexity of this procedure is $O(n \log n)$.

Let us now consider a special case of BQP01 when Q is of rank one (equivalently a special case of UBLP(1)) of the form

$$\begin{aligned} \text{BQP01}(1,0): \quad & \text{Maximize } (a_0 + ax)(b_0 + by) + cx + dy \\ & \text{Subject to: } x \in [0, 1]^m, y \in [0, 1]^n, \end{aligned}$$

where either $c = 0$ or $d = 0$. We now show that this problem can be solved more efficiently.

Theorem 9. *An optimal solution to BQP01(1,0) can be identified in $O(n)$ time.*

Proof. Suppose $d = 0$. Consider the following solutions

$$y_j^0 = \begin{cases} 1 & \text{if } b_j > 0 \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad y_j^* = \begin{cases} 1 & \text{if } b_j < 0 \\ 0 & \text{otherwise} \end{cases}.$$

Note that $y^0 = (y_1^0, y_2^0, \dots, y_n^0)$ maximizes $b_0 + by$ and $y^* = (y_1^*, y_2^*, \dots, y_n^*)$ minimizes $b_0 + by$ with respective objective function values, say, λ^0 and λ^* . Consider the function

$$L(\lambda) = \text{Maximum}_{x \in \{0,1\}^m} \{(a_0 + ax)\lambda + cx\}.$$

$L(\lambda)$ is a piecewise linear convex function of λ . As λ varies in the interval $[\lambda^*, \lambda^0]$, $L(\lambda)$ traces the best objective function values for BQP01(1,0) for all possible solution vectors y . Thus, BQP01(1,0) reduces to maximizing $L(\lambda)$ for $\lambda \in [\lambda^*, \lambda^0]$. Convexity of $L(\lambda)$ guarantees that its maximum is attained when $\lambda = \lambda^0$ or λ^* , i.e., when $y = y^0$ or $y = y^*$. But when $y = y^0$, an optimal x^0 is given by

$$x_i^0 = \begin{cases} 1 & \text{if } a_i \lambda^0 + c_i > 0 \\ 0 & \text{otherwise} \end{cases}.$$

Similarly, when $y = y^*$ an optimal solution x^* is given by

$$x_i^* = \begin{cases} 1 & \text{if } a_i \lambda^* + c_i > 0 \\ 0 & \text{otherwise} \end{cases}.$$

Thus, the best solution amongst (x^0, y^0) and (x^*, y^*) solves BQP01(1,0). For $c = 0$, the proof is similar. $\square$

We now show that our algorithm for solving BQP01 when rank of Q is fixed can be used to solve QP01 when Q' is positive semidefinite. Let $QP01(PSD, Q', c', n)$ denotes an instance of the QP01 on n variables on data Q', c' where Q' is positive semidefinite with fixed rank and let $BQP01(Q', c', n)$ denote an instance of BQP01 where $m = n$, $Q = Q'$ and $c = d = \frac{1}{2}c'$. Then we have,

Theorem 10. *If x^* is an optimal solution to $QP01(PSD, Q', c', n)$ and (x^0, y^0) be an optimal solution to $BQP01(Q', c', n)$. Then both x^0 and y^0 are optimal solutions to $QP01(PSD, Q', c', n)$ and (x^*, x^*) is an optimal solution to $BQP01(Q', c', n)$.*

The proof of the above theorem follows from a corresponding result from bilinear programming [17] and the equivalence between BQP01 and UBLP. An interesting consequence of this theorem is that our polynomial time algorithm for the case when rank of Q is fixed can be used to solve $QP01(PSD, Q', c', n)$ yielding a new alternative to the algorithms discussed in [1, 10, 16, 24] with the additional flexibility that c' could be arbitrary.

In the case of QP01, if $q_{ij} = a_i + b_j$ for $i \in I$, $j \in J$, the resulting problem is trivial. However, for BQP01, the condition $q_{ij} = a_i + b_j$ for $i \in I$, $j \in J$ yields an interesting polynomially solvable case. Note that when $q_{ij} = a_i + b_j$, rank of Q is at most 2 and hence can be solved in polynomial time. We now present an $O(mn \log n)$ algorithm to solve this problem.

Theorem 11. *If $q_{ij} = a_i + b_j$ for $i \in I$, $j \in J$, then BQP01 can be solved in $O(mn \log n)$.*

Proof. For any feasible solution x, y ,

$$\begin{aligned} f(x, y) &= x^T Q y + c x + d y + c_0 \\ &= \sum_{i=1}^m a_i x_i \sum_{j=1}^n y_j + \sum_{j=1}^n b_j y_j \sum_{i=1}^m x_i + \sum_{i=1}^m c_i x_i + \sum_{j=1}^n d_j y_j + c_0. \end{aligned}$$

Let $\sum_{j=1}^n y_j = K$ and $\sum_{i=1}^m x_i = L$, where K and L two parameters. Then,

$$f(x, y) = \sum_{i=1}^m (K a_i + c_i) x_i + \sum_{j=1}^n (L b_j + d_j) y_j + c_0.$$

Consider the optimization problem

$$\begin{aligned} \text{ILP}(K, L): \quad & \text{Maximize } \sum_{i=1}^m (K a_i + c_i) x_i + \sum_{j=1}^n (L b_j + d_j) y_j + c_0 \\ & \text{Subject to } \sum_{j=1}^n y_j = K \text{ and } \sum_{i=1}^m x_i = L \end{aligned}$$

For $K = 0, 1, \dots, n$, let α^K be a permutation of size m such that $K a_{\alpha^K(i)} + c_{\alpha^K(i)} \geq K a_{\alpha^K(i+1)} + c_{\alpha^K(i+1)}$, $i = 0, 1, \dots, m-1$. For $L = 0, 1, \dots, m$, let β^L be a permutation

of size n such that $Lb_{\beta^L(j)} + d_{\beta^L(j)} \geq Lb_{\beta^L(j+1)} + d_{\beta^L(j+1)}$, $j = 1, 2, \dots, n-1$. Observe that, for fixed K and L , the optimal $x = x^{K,L}$ can be obtained by setting $x_i = 1$ for $i = \alpha^K(1), \alpha^K(2), \dots, \alpha^K(L)$ and $x_i = 0$ for the rest of indices. Similarly, the optimal $y = y^{K,L}$ can be obtained by setting $y_j = 1$ for $j = \beta^L(1), \beta^L(2), \dots, \beta^L(K)$ and $y_j = 0$ for the rest of indices.

Let $f_1^{K,L} = \sum_{i=1}^m (Ka_i + c_i)x_i^{K,L}$ and $f_2^{K,L} = \sum_{j=1}^n (Lb_j + d_j)y_j^{K,L}$. Assume that we know the value of $f_1^{K,L}$ for some K and L . Then we can calculate $f_1^{K,L+1}$ in $O(1)$ time as:

$$f_1^{K,L+1} = f_1^{K,L} + Ka_{\alpha^K(L+1)} + c_{\alpha^K(L+1)}.$$

Observe also that $f_1^{K,0} = 0$. Hence, for a fixed K , we can calculate $f_1^{K,L}$ for each $L \in \{0, 1, \dots, m\}$ in $O(m)$ time. Similarly, for a fixed L , we can calculate $f_2^{K,L}$ for each $K = 0, 1, \dots, n$ in $O(n)$ time. Let (K^0, L^0) be the values of K and L that maximize $f_1^{K,L} + f_2^{K,L}$. Then the optimal solution of the BQP01 is $(x^{K^0,L^0}, y^{K^0,L^0})$, and it can be obtained in $O(nm \log m + mn \log n + mn) = O(mn \log n)$ time. $\square$

An integer linear programming formulation of BQP01 can be obtained using standard linearization techniques [14] as follows:

$$\text{ILP: Maximize } \sum_{i=1}^m \sum_{j=1}^n q_{ij} z_{ij} + \sum_{i=1}^m c_i x_i + \sum_{j=1}^n d_j y_j \quad (17)$$

$$\text{Subject to } z_{ij} - x_i \leq 0, i \in I, j \in J \quad (18)$$

$$z_{ij} - y_j \leq 0, i \in I, j \in J \quad (19)$$

$$z_{ij} - x_i - y_j \geq -1, (i, j) \in S, \quad (20)$$

$$x_i \in \{0, 1\} \text{ for } i \in I, y_j \in \{0, 1\} \text{ for } j \in J, z_{ij} \in \{0, 1\} \text{ for } (i, j) \in I \times J, \quad (21)$$

where $S = \{ij : q_{ij} < 0\}$. Note that each constraint (20) forces the variable z_{ij} to take value 1 if both x_i and y_j are one. We introduce such a constraint only when $q_{ij} < 0$. If $q_{ij} > 0$, then, since the objective function is being maximized, z_{ij} will be forced to take the value 1 whenever both x_i and y_j are 1 and hence explicit constraint in this case is unnecessary. If $q_{ij} = 0$, the value of z_{ij} is irrelevant. This leads to another polynomially solvable case.

Theorem 12. *Let $|\{ij : q_{ij} < 0\}| = p$. Then BQP01 can be solved in polynomial time if $p = O(\log n)$ and it is NP-hard if $p = \sqrt[k]{n}$.*

Proof. Let $p = O(\log n)$. Deleting constraints (20) from ILP results in a totally unimodular system which can be solved in polynomial time. There are at most $3p$ binary variables associated with equation (20). Thus, there are at most 2^{3p} possible combinations of values of these variables. Substituting each such combination in ILP results in the elimination of the constraints and variables corresponding (20) and the resulting system will be totally unimodular and have at most two variables per inequality. Thus the reduced ILP can be solved in polynomial time [25]. Thus, ILP can be solved by choosing the best solution amongst the solutions of these reduced ILPs. Since the total number of reduced ILPs is

at most 2^{3p} and $p = O(\log n)$, the number of reduced ILP considered is $O((n)^{3k})$ for some constant k and the result of the first part follows.

For the second part, the proof follows by reducing a general BQP01 to a BQP01 satisfying the conditions of the theorem. This can be achieved by increasing the number of columns of Q to sufficiently large, yet polynomial for fixed k and filling these columns with entries 0. $\square$

It may be noted that Theorem 12 allows arbitrary c and d . The special case of Theorem 12 where $q_{ij} \geq 0$ for all (i, j) has been reported by many authors [27] in the context of QP01. The general version discussed in Theorem 12 for BQP01 extends to QP01 as well.

4 Conclusion

In this paper we studied the problem BQP01 which generalizes QP01, a well studied combinatorial optimization problem. BQP01 is known to be MAX SNP hard. Several interesting polynomially solvable special cases of the problem are identified. In particular, we showed that when rank of the matrix Q is fixed, BQP01 can be solved in polynomial time and improved complexity results are provided for the rank one case and a special case when rank of Q is at most two. By restricting $m = O(\log n)$ or by restricting the number of negative entries of Q to $O(\log n)$, we obtained additional polynomially solvable cases. If $m = \sqrt[k]{n}$ BQP01 is MAX SNP hard and a similar result is obtained if the number of negative entries in Q is $O(\sqrt[k]{n})$. It would be interesting to explore the complexity of the problem when the problem size or data restrictions fall in between these extreme cases. Exploiting the algorithms for our polynomially solvable special cases, efficient exact and heuristic algorithms may be obtained to solve BQP01. This is a topic for further investigation.

References

- [1] K. Allemand, K. Fukuda, Th. M. Liebling, and E. Steiner. A polynomial case of unconstrained zero-one quadratic optimization. *Mathematical Programming, Ser. A*, 91 (2001) 49-52.
- [2] N. Alon and A. Naor, Approximating the cut-norm via Grothendieck's inequality, *SIAM Journal of Computing* 35 (2006) 787-803.
- [3] M. Altman, Bilinear programming, *Bullentin de l' Académie Polonaise des Sciences* 16 (1968) 741-746.
- [4] C. Ambühl, M. Mastrolilli, and O. Svensson, Inapproximability results for maximum edge biclique, minimum linear arrangement, and sparsest cut, *SIAM Journal of Computing* 40 (2011) 567-596.
- [5] E. Boros and P. L. Hammer, The max-cut problem and quadratic 01 optimization; polyhedral aspects, relaxations and bounds, *Annals of Operations Research* 33 (1991) 151-180.

- [6] E. Boros, P.L. Hammer, and G. Tavares. Local search heuristics for Quadratic Unconstrained Binary Optimization. *Journal of Heuristics* 13 (2007) 99-132.
- [7] E. Cella, B. Klinz, and C. Meyer, Polynomially solvable cases of the constant rank unconstrained quadratic 0-1 programming problem, *Journal of Combinatorial Optimization* 12 (2006) 187-215.
- [8] W.-C. Chang, S. Vakati1, R. Krause, and O. Eulenstein, Exploring biological interaction networks with tailored weighted quasi-bicliques, *BMC Bioinformatics* 13(Suppl 10):S16, 2012.
- [9] M. Charikar and A. Wirth, Maximizing quadratic programs: extending Grothendieck's inequality, FOCS 2004, 54-60.
- [10] J.-A. Ferrez, K. Fukuda, and Th.M. Liebling, Solving the fixed rank convex quadratic maximization in binary variables by a parallel zonotope construction algorithm. *European Journal of Operational Research*, 166 (2005) 35–50.
- [11] N. Gillis and F. Glineur, Low-rank matrix approximation with weights or missing data is NP-hard, *SIAM Journal of Matrix Analysis and Applications* 32 (2011) 1149-1165.
- [12] M. X. Goemans and D. P. Williamson, Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming, *Journal of the Association for Computing Machinery* 42 (1995) 1115-1145.
- [13] P. L. Hammer, P. Hansen, P. Pardalos, and D. J. Rader, JR, Maximizing the product of two linear functions in 0-1 variables, *Optimization* 51 (2002) 511-537.
- [14] P. L. Ivănescu, Some network flow problems solved using pseudo-boolean programming, *Operations Research* 13 (1965) 388-399.
- [15] T. Gal, *Postoptimal analysis, parametric programming, and related topics*, Walter de Gruyter, Berlin, 1995.
- [16] G. N. Karystinos and A. P. Liavas, Efficient computation of the binary vector that maximizes a rank-deficient quadratic form, *IEEE Transactions on information theory*, 56 (2010) 3581 – 3593.
- [17] H. Kono, Maximization of a convex quadratic function under linear constraints, *Mathematical programming* 11 (1976) 117-127.
- [18] H. Konno, T. Kuno, and Y. Yajima, Parametric simplex algorithms for a class of NP-complete problems whose average number of steps is polynomial, *Computational optimization and applications* 1 (1992) 227-239.
- [19] M. Koyutürk, A. Grama, and N. Ramakrishnan, Compression, clustering, and pattern discovery in very high-dimensional discrete-attribute data sets, *IEEE Transactions on Knowledge and Data Engineering*, 17 (2005) 447-461.

- [20] M. Koyutürk, A. Grama, and N. Ramakrishnan, Nonorthogonal Decomposition of Binary Matrices for Bounded-Error Data Compression and Analysis, *ACM Transactions on Mathematical Software* 32 (2006) 33-69.
- [21] D. Li, X. Sun, S. Gu, J. Gao, and C. Liu, Polynomially solvable cases of binary quadratic programs, in Optimization and Optimal Control, A. Chinchuluun et al. (eds), Springer Optimization and its Applications 39, 2010.
- [22] Z. Lü, F. Glover, and J.-K. Hao, A hybrid metaheuristic approach to solving the UBQP problem, *European Journal of Operational Research* 207 (2010) 1254-1262.
- [23] H. Lu, J. Vaidya, V. Atluri, H. Shin, and L. Jiang, Weighted rank-one binary matrix factorization
- [24] G. Manglani and A. K. Chaturvedi, Application of computational geometry to multiuser detection in CDMA, *IEEE Transactions on Commun.*, 54 (2006) 204-207.
- [25] N. Megiddo, Towards a genuinely polynomial algorithm for linear programming, *SIAM Journal of Computing* 12 (1983) 347-353.
- [26] K. G. Murty, *Linear programming*, John Wiley & Sons, New York, 1983.
- [27] J.C.Picard and H.D.Ratliff, Minimum cuts and related problems, *Networks* 5 (1974)357-370.
- [28] P. Raghavendra and D. Steurer, Towards computing the Grothendieck constant. In Proceedings of the twentieth Annual ACM-SIAM Symposium on Discrete Algorithms (SODA '09). Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 525-534, 2009.
- [29] B.-H. Shen, S. Ji, and J. Ye, Mining discrete patterns via binary matrix factorization, In KDD'09: Proceedings of the 15th ACM SIGKDD international conference on Knowledge discovery and data mining, New York, USA, (2009) 757-766.
- [30] J. Tan. Inapproximability of maximum weighted edge biclique and its applications. In Proceedings of the 5th international conference on Theory and applications of models of computation (TAMC'08), Manindra Agrawal, Dingzhu Du, Zhenhua Duan, and Angsheng Li (Eds.). Springer-Verlag, Berlin, Heidelberg, 282-293, 2008.
- [31] A. Tanay, R. Sharan, and R. Shamir, Discovering statistically significant biclusters in gene expression data, *Bioinformatics* 18 (2002) Suppl. 1, S136-S144.
- [32] Y. Yajima and H. Konno, Efficient algorithms for solving rank two and rank three bilinear programming problems, *Journal of global optimization* 1 (1991) 155-171.